

Study of the mechanism of reactions catalyzed by complexes of cytochrome C with cardiolipin

Levchenko I.N.^{1*}, Volodyaev I.V.², Boyarchenkov A.S.³

¹ Pirogov Russian National Research Medical University, Moscow, Russia

² Lomonosov Moscow State University, Moscow, Russia

³ Ural Federal University named after the first President of Russia B.N. Yeltsin, Ekaterinburg, Russia

* irnlevchenko@yandex.ru

Key words: cytochrome C, peroxidases, nanomachine, peroxidases, mathematical modeling

Motivation and Aim: When studying objects of the living world, we often encounter phenomena that are well described by algorithms from the category of nanomachines. Models of this class have existed for more than 50 years and effectively help mankind in understanding many phenomena and solving practical problems. Thus, in many cases, simplified models of nanomachines allow a fairly good understanding of complex multi-component biological processes. For instance, the complex of cytochrome C with the phospholipid cardiolipin, interpreted as a nanomachine, quite well describes one of the key nodes in triggering apoptosis – interaction of cytochrome C with the inner membrane of mitochondria, activating its peroxidase activity and further release into the cytoplasm. Cytochrome C is a heme-containing protein that promotes electron transfer between mitochondrial respiratory complexes 3 and 4, and participates in triggering apoptosis. As shown [1], cytochrome C forms a complex with the membrane phospholipid cardiolipin, which activates its peroxidase activity and initiates lipid peroxidation in the mitochondrial membranes. This breaks the membrane barrier properties, releasing cytochrome C into the cytoplasm, where it binds to the protein Apaf-1, forming apoptosomes and triggering apoptosis. The general scheme of its peroxidase action is presented in Fig. 1.




Fig. 1. General reaction scheme of mammalian peroxidases

Regarded as a nanomachine, the Cytochrome C / cardiolipin complex is especially valuable as a trigger, switching between normal cell functioning and apoptosis

depending on its peroxidase activity. Generally speaking, the latter should be measured by the rate of free radical generation, but due to their extremely high reactivity, it cannot be done directly. Thus, it is estimated by the intensity of oxychemiluminescence, accompanying further reactions of the generated free radicals, and has to be done in dynamics – which demands the use of the so-called kinetic chemiluminescence (see [5]). *Methods and Algorithms:* Having the above experimental data available, in this work, we describe the mathematical model, creating an automated solution to the direct and inverse problems to study the mechanism of cytochrome C participation in apoptosis. Importantly, in this work we consider only the control process, which is carried out by changing the protein/lipid structural ratio and membrane viscosity.

The reaction rates were expressed as the product of the concentration of the reactants and the corresponding rate constant (see Fig. 1). The rate constants were selected manually. The initially formed system of differential equations for reaction rates was solved by the Euler method.

Further, specific optimization was proposed for solving the inverse problem. Several algorithms, each of which could serve as the basis for optimizing the selection process both for the rate constants and for the concentrations of the given substances, were progressively compared with the experiment, with the best one finally selected.

Conclusion: A program for the automatic determination of the rate constants of chemical reactions was created and debugged, corresponding to the formulation of the inverse problem.

Unsatisfactory solutions were screened, and the best model selected.

The rate constants of the reactions proceeding in the system for measuring the activity of the luminescence intensity were determined.

Acknowledgements: We would like to express our gratitude to the supervisors for their valuable contribution to the planning of the study, namely, Academician, Professor Yuri Andreevich Vladimirov, Faculty of Fundamental Medicine, Moscow State University and Professor Anatoly Nikolaevich Osipov, Faculty of Medicine and Biology, Pirogov Russian National Research Medical University.

References

1. Vladimirov Y.A. et al. Mechanism of activation of cytochrome C peroxidase activity by cardiolipin. *Biochemistry (Mosc)*. 2006;71(9):989-97.
2. Cortese J.D., Voglino A.L., Hackenbrock C.R. The ionic strength of the intermembrane space of intact mitochondria is not affected by the pH or volume of the intermembrane space. *Biochim Biophys Acta*. 1992;1100(2):189-197.
3. Nantes I.L. et al. Modifications in heme iron of free and vesicle bound cytochrome C by tert-butyl hydroperoxide: a magnetic circular dichroism and electron paramagnetic resonance investigation. *Free Radic Biol Med*. 2000;28(5):786-796.
4. Dawson J.H. Probing structure-function relations in heme-containing oxygenases and peroxidases. *Science*. 1988;240(4851):433-439.
5. Furtmuller P.G. et al. Kinetics of interconversion of redox intermediates of lactoperoxidase, eosinophil peroxidase and myeloperoxidase. *Jpn J Infect Dis*. 2004;57(5):S30-S31.
6. Weng L.C., Baker G.M. Reaction of hydrogen peroxide with the rapid form of resting cytochrome oxidase. *Biochemistry*. 1991;30(23):5727-5733.
7. Kagan V.E. et al. A role for oxidative stress in apoptosis: oxidation and externalization of phosphatidylserine is required for macrophage clearance of cells undergoing Fas-mediated apoptosis. *J Immunol*. 2002;169(1):487-499.
8. Kagan V.E. et al. Cytochrome C acts as a cardiolipin oxygenase required for release of proapoptotic factors. *Nat Chem Biol*. 2005;1(4):223-232.